

## PROJECT PROJECT



Fentanyl Education And Naloxone Training



#### Please take our anonymous survey

How much do you know about Fentanyl?



# DON'T TRY. DON'T DIE.



3/16/2002 - 9/25/2021



#### As reported by the CDC:

 There were an estimated 100,306 drug overdose deaths in the US for the 12-month period ending April 2021.

 The CDC indicated that more than 107,000 Americans died from a drug overdose in the 12months ending in January 2022.

100,306

**April** 

107,000

January



Overall, 75% involved at least one

opioid, with 71,450 (66.5%) involving

synthetic opioids, primarily illicitly

manufactured Fentanyl or Fentanyl

analogs.

71,450

66.5% Synthetic Opioids

PROJECT O D

### **The Numbers**



- The number of adolescent (ages 14-18) overdose deaths rose by 114% between 2019 and 2021 with fentanyl accounting for 77% of adolescent overdose deaths
- Among adolescent overdose decedents (ages 10 to 19) only 35% of them had previous opioid use history, yet 90% of deaths were opioid related, and 24.5% had evidence of counterfeit pill involvement.

114%

2019-2021

90%

Opioid Related

## FAMOUS FENTANYL FATHERS

#### **ERIC BOLLING**

#### MASTER P ROBERT DE NIRO

Eric Bolling's Son 'Wanted To Get Rid Of The Pain' Before Tragic OD Death, Pal Claims

Master P posts emotional message after daughter Tytyana's death: 'Life is too short'

Grieving Robert De Niro Spotted for First Time Since Death of 19-year-old Grandson Leandro, Visits Funeral Home With Daughter Drena







"Fentanyl" is mentioned 175,000 a day in broadcast and published articles.\*

\*LexisNexis.com

#### **The Numbers**



- Overdoses from opioids increased to
  80,590 from 76,000 1 year prior.
- Stimulant overdose deaths are also on the rise, with approximately 33,128 (30.8%) deaths involving methamphetamine and 24,751 (23%) involving cocaine.

4,590

Overdose Increase

30.8%

Methamphetamine

23%

Cocaine

#### Who - Who We Are



#### Fentanyl Fathers

A non-profit working to educate kids and warn communities about the dangers of fentanyl and counterfeit pills through the lens of heartbroken parents. Our goal is to spare families from waking up to the same grief we share in every day.

#### Mission:

To establish a world where no adolescent succumbs to fentanyl poisoning or overdose, by raising awareness about the dangers of fentanyl, counterfeit pills, and effective overdose response measures.

#### What - We Will Do Today

#### Goals for this presentation:

- Educate and raise awareness on the dangers of Fentanyl and counterfeit pills
- Train to administer Nasal Naloxone in the event of an opiate/opioid drug overdose
- Raise awareness about how Fentanyl poisoning and overdose death affects families





#### What - We Will Do Today



#### Objectives:

- Be able to recognize and respond to an opiate drug overdose
- Increase the perceived threat of experiencing or witnessing an overdose
- Raise awareness about the potential consequences of drug experimentation
- Learn how to administer Nasal Naloxone
- Connect people to important resources and eliminate access barriers
- Understand the benefits of drug abstinence


#### Why – Why We Do It



- To reduce stigma and misinformation through education to our community
- To have Naloxone more readily available to minimize the number of lives lost to overdose and accidental poisonings
- To raise community capacity for recognizing and responding to an emerging public health threat that is killing more people today than ever before

#### Why – Why We Do It

- To ensure that no family or community experiences the grief of losing a loved one to an overdose or drug poisoning without warning
- To warn kids about the dangers of the unpredictable illicit and counterfeit pill drug supply ever-present in America today FATHERS
- Ultimately, we do this to save lives
- We lost our kids to this. Saving a life in their name makes their life MATTER.




#### The Impact

Deaths By Category	Deaths
2022 Drunk Driving +	10,360
9/11 Attack +	2,977
Covid Deaths (18-29) +	6,749
2022 Childhood Cancer +	1,600
Vietnam War Deaths	58,220
Global Plane Crashes since 2006	14,572
Total	94,478












**Jack McCarthy** 



### **Jack McCarthy**





### **Jack McCarthy**







Potentially Lethal Dose of Fentanyl
























#### FAKE PRESCRIPTION PILLS • WIDELY AVAILABLE • INCREASINGLY LETHAL

DEA LAB TESTING REVEALS THAT

7 OUT OF EVERY 10 PILLS

WITH FENTANYL CONTAIN A POTENTIALLY

## LETHAL DOSE

COUNTERFEIT PILLS OFTEN CONTAIN FENTANYL AND ARE MORE LETHAL THAN EVER BEFORE



# FATHERS

### **Counterfeit Pills**

- Mass produced by criminal drug networks
- Easy to purchase, widely available
- Often contain fentanyl or methamphetamine
- Sold on social media and e-commerce platforms (anyone can purchase with a smartphone, including minors)



#### **Counterfeit Pills**

- Fentanyl is a synthetic opioid that is
  50x more potent than heroin and 100x
  more potent than morphine
- In 2022, the DEA found 6 out of 10 fake pills contain a lethal dose of fentanyl
  - 2 mg is enough to kill you
  - There are 2000 mg in a sugar packet by comparison



#### **Counterfeit Pills**

- These pills are being sold on social media to look like Oxycontin, Percocet, Vicodin, Xanax, and stimulants like Adderall
- Assume every pill you purchase today WILL KILL YOU
- Fentanyl powder is also being added to street drug supplies like cocaine, heroin, methamphetamine




#### What Is An Opioid Overdose?



Opioids Attaching To Receptors

The brain has many, many receptors for opioids. An overdose occurs when too much of an opioid, such as heroin or oxycodone, fits in too many receptors slowing and then stopping the breathing.



#### Naloxone Is Effective

The life-saving antidote works to reverse opiate overdoses from the following drugs:

#### Heroin

- Methadone
- Morphine
- Vicodin/Norco
- Percocet
- Codeine

#### Lortab

- Tramadol
- Oxymorphone
- Hydrocodone
- Oxycodone
- Fentanyl






#### **Naloxone In The Brain**

Naloxone Reversing An Overdose

Naloxone has a stronger affinity to the opioid receptors than opioids, such as heroin or oxycodone, so it knocks the opioids off the receptors for a short time (30-90 minutes). This allows the person to breathe again and reverse the overdose.



#### **Naloxone In Action**



- Works on ANY opiate ANY time
- Reverses opiate effects- sedation and respiratory depression
- Causes sudden withdrawal an unpleasant experience
- No psychoactive effects not addictive
- Routinely used by EMS (often in larger doses)



There will be no effect on the individual if an opiate is not present

#### Signs And Symptoms Of Opioid USE

- Nodding out/Falling asleep
- Pinpoint pupils
- May see paraphernalia laying around
- Alarming or unusual changes in behavior







#### **Pupil Variations**

# FATHERS

**Opiate vs. Stimulant Pupil Changes** 


Normal



Dilated



Pinpoint



#### Signs And Symptoms Of Opioid OVERDOSE

- NO breathing or shallow breathing
- Limp
- Turning pale or blue especially in lips and fingernails
- Gurgling or snoring noises





#### Nasal Narcan Save A Life Kit

#### **Each Save A Life Kit Contains:**

- 2 Doses of 4mg Nasal Narcan
- Sterile gloves
- Reference card
- Community Resource Guide
- Sober Support Unit information
- Fentanyl Testing Strip (as requested)



PROJECT PROJECT

Step 1: Call 9-1-1

- Request EMS or fire department
- Try to put gloves on to protect yourself

Law Act 307 Public Acts of 2016

- (7) As used in this section:
- (b) "Seeks medical assistance" means reporting a drug overdose or other medical emergency to law enforcement, the 9-1-1 system, a poison control center, or medical provider, or assisting someone in reporting a drug overdose or other medical emergency.





Step 2: Assess Situation

Do not disturb the area

Check consciousness









### Step 3: Give Nasal Narcan

- Remove 1 4mg dose of the Nasal Narcan
- Tilt individual's head back
- Dispense dose as directed into 1 nostril











Where to store our FREE Naloxone. White AED Unit *or* Red Stand Alone.

PROJECT PIOID







## Step 4: Possible 2<sup>nd</sup> Dose

- Wait 3 minutes
- If no change in victim's condition, administer the second dose in the opposite nostril



PROJECT PIOID

### Step 5: Withdrawal

- Place individual on their side in the recovery position (left side)
- Possible vomiting, diarrhea, and/or sweating
- Possible agitated behavior







Don't Try. Don't Die.

Please complete the survey!

PROJECT PROJECT





## CONTACT

Greg Swan | 248 667 2722 | Bob Minotti | 386 801 4884 Greg@FentanylFathers.org. | Bob.Minotti@FentanylFathers.org.

Website: <u>FentanylFathers.org</u>

HQ: New Smyrna Beach, Florida



### MEDIA COVERAGE

Highlights 279 Die CBS Alaska ABC Alaska

Alaska Assembly Dr. Phil Dr. Phil 2 FOX Alaska PSA

NBC Ohio NBC Ohio 2 NBC Ohio 3 Detroit Free Press Op-Ed

Detroit Free Press 2 Ohio Radio ABC Ohio Michigan Assembly NBC

ALASKA|CALIFORNIA|FLORIDA|MICHIGAN|OHIO|TEXAS|WASHINGTON